

Acknowledgments

Research into the systematics of *Melaleuca* in Canberra, Australia, was initiated by Bryan Barlow to whom credit is given for leading the early studies and demonstrating what a rewarding genus it is for scientific investigations. In the early 1980s, one of us (LAC) began working on the taxonomy of a group of species then considered to belong in *Melaleuca* but now recognised as *Asteromyrtus*, a genus allied to *Agonis*. Subsequently, LAC extended his research into the *M. scabra* group of species, into other parts of the genus and into other concepts within the *Melaleuca* complex. A great many people have assisted these studies through their provision of specimens, companionship and assistance in the field and/or herbarium, hospitality, information on the existence of unusual populations, and so on. These people are too numerous to name individually but we thank them here for their generosity and willingness to go out of their way to help. Special thanks, however, are due to colleagues then and/or currently on the staff of the Australian National Herbarium, notably Bryan Barlow, Kirsten Cowley, Brendan Lepschi, Julie Matarczyk and Frances Quinn.

When this book was conceived, it was evident that no information on essential oils was available for over 90 species, despite the authors, wherever possible, having made a point of collecting foliage samples for distillation and analysis over the years. Funding was provided by the Rural Industries Research and Development Corporation (RIRDC) to enable a dedicated collecting trip to be made in south-western Western Australia in 2010, during which many of the outstanding species were collected for oil analysis. This trip was undertaken by one of us (JCD) and David Lea whose help on this trip and in undertaking several additional collections is gratefully acknowledged. Requests have also been made of people and organisations in a position, potentially, to assist us to obtain samples. In most cases, these requests resulted in the collection of samples, thus enabling analyses to be prepared of most of the missing taxa. These people and/or organisations are numerous and, in some cases, the samples that we received were collected as a result of our request having been passed to other parties. To attempt to name each person individually would result in an incomplete list but we acknowledge with sincere thanks the goodwill and assistance of those who provided samples.

The majority of the photographs in the ‘Species accounts’ (Chapter 7) were made available to us by the late Ivan Holliday. Thanks are due to the many other people who have contributed photographs; they are named in the photograph credits below. We thank Anto Rimbawanto, Le Dinh Kha and Stephen Midgley for providing information on the utilisation of melaleucas in Indonesia, Vietnam and China, respectively. Peter White provided information on the use of *M. viminea* sticks in the lobster-fishing and vegetable-growing industries in Western Australia. Carl Davies and Nunzio Knerr, both with CSIRO, deserve special thanks. Carl digitised transparencies and prepared the digital files for publication, while Nunzio prepared the distribution maps and generated the species-richness and endemism maps. We thank Christian Doran for his editorial work on the photographs and figures presented in the draft introductory chapters. The passage on the use of melaleucas in bonsai was based on notes written for us by Roger Hnatiuk; it is unfortunate that space limitations prevented further elaboration of this interesting topic. Murray Fagg, Australian National Botanic Gardens, generously permitted us to use his illustrations of floral details of typical *Melaleuca* species (Figures 3 and 4).

We extend special thanks to John Turnbull, Bill Foley, Ian Southwell and Kirsten Cowley for their helpful comments on a draft of this manuscript.

This volume is dedicated to the memories of Ivan Grenfell Holliday (1926–2010) sportsman, plant photographer and author, and Robert John Goldsack (1941–2009) chemist and botanist. Ivan Holliday was keenly interested in *Melaleuca* and produced a volume (with two editions) on the genus. He accompanied Lyn Craven and Brendan Lepschi on a key collecting mission to south-western Western Australia in 1994. Bob Goldsack, who died tragically in 2009, was both a botanist and chemist who greatly contributed for over 20 years to the work on the essential oils of the melaleucas. His tireless work on the chemistry of this genus, as well as his fresh outlook on ‘life, the universe and everything’ is greatly missed by all who came into contact with him.

Photograph credits

Chapters 1–6: In addition to the authors’ contributions, the following persons or organisations provided some or all of the images in the following figures: Figure (Fig.) 1, Christine Doran; Fig. 2, Australian Tree Seed Centre (ATSC); Figs 3 & 4, Murray Fagg; Figs 5, 6, 7 & 8, Liliana Baskorowati; Fig. 11A, Ivan Holliday; Fig. 11B, Wayne O’Sullivan; Fig. 11C, Nathan McQuoid; Fig. 12, ATSC & Forest Science Institute of Vietnam (FSIV); Fig. 13, ATSC & David Lea; Fig. 15A, Frank Zich; Fig. 15B, Murray Fagg; Fig. 16A, Roger Hnatiuk; Fig. 16B, David Sweet; Fig. 17A, Murray Fagg; Fig. 18B, C, D, E, F, Murray Fagg; Fig. 19, FSIV; Fig. 20, Mary Webb; Fig. 23, ATSC; Fig. 26A, Anto Rimbaunto; Fig. 27, Peter Entwistle; Fig. 28, Gary Baker & Liliana Baskorowati.

Species accounts (Chapter 7): The following people and/or organisations provided images for the listed species: David Albrecht: *interioris*; Australian National Botanic

Gardens: *biconvexa* (photo: D. Greig); Liliana Baskorowati: *dissitiflora*; Don Bellairs (Ivan Holliday Image Library): *hollidayi*; Kym Brennan: *citrolens*, *dealbata*, *ferruginea*, *sericea*, *stipitata*; Centre for Australian National Biodiversity Research: *apostiba* (photo: J. Connors); L. Carman (Ivan Holliday Image Library): *tinkeri*; Colin Cornford (Ivan Holliday Image Library): *arcana*, *cornucopiae*, *foliolosa*; Ian Cowie: *triumphalis*; Gil Craig: *johsonii*, *papillosa*, *pau-periflora*, *pomphostoma*, *ulicoides*; Lyn Craven: *atroviridis*, *megalongensis*, *nematophylla*, *punicea*, *uncinata*, *vinnula*, *zeteticorum*; Mike Crisp: *agathosmoides*, *amydra*, *camptoclada*, *glaucha*, *grieveana*, *halophila*, *idana*, *pritzelii*, *sabrina*, *sciotostyla*, *serpentina*; Christine Doran: *armillaris*; Bill Dowling: *nanophylla*; Murray Fagg, Australian National Botanic Gardens: *acacioides*, *alsophila*, *argentea*, *blaerii-folia*, *brachyandra*, *cajuputi*, *citrina*, *comboynensis*, *decora*, *densispicata*, *ericifolia*, *faucicola*, *flammea*, *flavovirens*, *formosa*, *globifera*, *groveana*, *leptospermoides*, *linariifolia*, *megacephala*, *micromera*, *minutifolia*, *montana*, *nervosa*, *nodosa*, *pachyphylla*, *paludicola*, *parvistaminea*, *phratra*, *podiocarpa*, *quinquenervia*, *recurva*, *salicina*, *shiressii*, *squamophloia*, *stenostachya*, *stypheleoides*, *subulata*, *tamariscina*, *thyoides*, *viminalis*, *violacea*, *virens*, *williamsii*; Andrew Ford: *borealis*, *monantha*, *sylvana*, *uxorum*; Mary Hoggart, Esperance Wildflower Society: *dempta*, *exuvia*, *fissurata*, *linguiformis*, *phoidophylla*, *plumea*, *sparsiflora*, *subalaris*; David Lea: *bisulcata*, *cliffortioides*, *genialis*, *haplantha*, *holosericea*, *protrusa*, *similis*, *urceolaris*; Glenn Leiper: *howeana*; Daniel & Irène Létocart: *dawsonii*, *gnidioides*, *pancheri*, *sphaerodendra*; Nathan McQuoid: *sophisma*; Neil Marriott: *wimmerensis*; Rhonda Melzer: *hemisticta*; Peter Olde: *deanei*; Wayne O’Sullivan: *osullivanii*, *scalena*, *teuthidoides*; Steven & Alison Pearson: *chisholmii*, *lazaridis*, *pearsonii*, *trichostachya*; Joel Plumb: *fluviaialis*; Anne Rick: *lecanantha*, *sculponeata*; Jean-Louis Ruiz: *brongniartii*, *buseana*; Garry & Nada Sankowsky: *polandii*, *pyramidalis*; John Simmons (Ivan Holliday Image Library): *pustulata*; Corin Urquhart: *linearifolia*; Peter White: *eurystoma*; Frank Zich: *stereophloia*, *zonalis*. Ivan Holliday provided all the remaining photographs in this chapter.

Copyright for the images in this volume is held by the person and/or entity or organisation named in the lists above.

Appendix 1 Oil types by species

Oil types found in the various *Melaleuca* species are based on a broad classification of the composition of the oils found. The basis for each oil type is described in the box below. Summaries of the oil composition of the species are provided in the ‘Species accounts’ (Chapter 7) and full details are accessible at <<http://aciar.gov.au/publication/MN156>>.

#	Species	Oil type
1	<i>M. acacioides</i>	J
2a	<i>M. acuminata</i> subsp. <i>acuminata</i>	A, L
2b	<i>M. acuminata</i> subsp. <i>websteri</i>	C, L

Oil compositions used in the table

Where percentage figures are mentioned, they are to be taken as only approximate values.

A	50–70% 1,8-cineole, <10% each of α -pinene, limonene, α -terpineol
B	20–50% terpinen-4-ol
C	major amounts of terpinolene
D	lemon-scented oils
E	aromatic oils (in the chemical sense)
F	major amounts of E-nerolidol
G	large amounts of α -pinene
H	significant amounts of β -diketones
J	other sesquiterpenic oils
K	significant amounts of linalool
L	other monoterpenoid oils
M	virtually no oil present
X/Y	oil compositions midway between X and Y type oils
(Z)	isolated occurrences of Z type oils but by no means the major oil type
-	no data
#	species number (notional)

#	Species	Oil type
3	<i>M. acutifolia</i>	A
4	<i>M. adenostyla</i>	A
5	<i>M. adnata</i>	L
6	<i>M. agathosmoides</i>	L
7	<i>M. alsophila</i>	G, D, B, A
8	<i>M. alternifolia</i>	A, B, C
9	<i>M. amydra</i>	A
10	<i>M. apodocephala</i>	J/L
11	<i>M. apostiba</i>	A
12	<i>M. araucariooides</i>	G
13	<i>M. arcana</i>	A, B
14	<i>M. argentea</i>	L, F
15a	<i>M. armillaris</i> subsp. <i>akineta</i>	A
15b	<i>M. armillaris</i> subsp. <i>armillaris</i>	A
16	<i>M. aspalathoides</i>	L
17	<i>M. atroviridis</i>	A
18	<i>M. barlowii</i>	L
19	<i>M. basicephala</i>	A
20	<i>M. beardii</i>	G
21	<i>M. biconvexa</i>	J
22	<i>M. bisulcata</i>	K
23	<i>M. blaeriifolia</i>	A
24	<i>M. boeophylla</i>	L

#	Species	Oil type
25	<i>M. borealis</i>	A
26	<i>M. brachyandra</i>	M
27	<i>M. bracteata</i>	E, (J)
28	<i>M. bracteosa</i>	J
29	<i>M. brevifolia</i>	L
30	<i>M. brevisepala</i>	J
31	<i>M. bromelioides</i>	A
32	<i>M. bronniartii</i>	G/L
33	<i>M. brophyi</i>	A
34	<i>M. buseana</i>	C/L
35	<i>M. caeca</i>	A
36a	<i>M. cajuputi</i> subsp. <i>cajuputi</i>	A, F
36b	<i>M. cajuputi</i> subsp. <i>cumingiana</i>	L
36c	<i>M. cajuputi</i> subsp. <i>platyphylla</i>	G, H
37	<i>M. calcicola</i>	B
38	<i>M. calothamnoides</i>	A

Oil compositions used in the table

Where percentage figures are mentioned, they are to be taken as only approximate values.

A	50–70% 1,8-cineole, <10% each of α -pinene, limonene, α -terpineol
B	20–50% terpinen-4-ol
C	major amounts of terpinolene
D	lemon-scented oils
E	aromatic oils (in the chemical sense)
F	major amounts of E-nerolidol
G	large amounts of α -pinene
H	significant amounts of β -diketones
J	other sesquiterpenic oils
K	significant amounts of linalool
L	other monoterpenoid oils
M	virtually no oil present
X/Y	oil compositions midway between X and Y type oils
(Z)	isolated occurrences of Z type oils but by no means the major oil type
-	no data
#	species number (notional)

#	Species	Oil type
39	<i>M. calycina</i>	L, E
40	<i>M. calyptroides</i>	A
41	<i>M. campanae</i>	L
42	<i>M. campyoclada</i>	A
43	<i>M. capitata</i>	G
44	<i>M. cardiophylla</i>	L
45	<i>M. carrii</i>	A, L
46	<i>M. cheelii</i>	G
47	<i>M. chisholmii</i>	A
48	<i>M. ciliosa</i>	K
49	<i>M. citrina</i>	A
50	<i>M. citrolens</i>	A, L, D, E
51	<i>M. clarksonii</i>	J
52	<i>M. clavifolia</i>	L
53	<i>M. cliffortioides</i>	J
54	<i>M. coccinea</i>	J
55	<i>M. comboynensis</i>	A
56	<i>M. concinna</i>	G/J
57	<i>M. concreta</i>	A/B
58	<i>M. condyllosa</i>	L
59	<i>M. conothamnoides</i>	L
60	<i>M. cordata</i>	L
61	<i>M. cornucopiae</i>	L
62	<i>M. croxfordiae</i>	A
63	<i>M. ctenoides</i>	A
64	<i>M. cucullata</i>	J
65	<i>M. cuticularis</i>	A
66	<i>M. dawsonii</i>	C/J
67	<i>M. dealbata</i>	J
68	<i>M. deanei</i>	G
69	<i>M. decora</i>	G, L
70	<i>M. decussata</i>	A
71	<i>M. delta</i>	A
72	<i>M. dempta</i>	L
73	<i>M. densa</i>	A
74	<i>M. densispicata</i>	A

#	Species	Oil type
75	<i>M. depauperata</i>	A
76	<i>M. depressa</i>	K
77	<i>M. dichroma</i>	A
78	<i>M. diosmatifolia</i>	A
79	<i>M. diosmifolia</i>	A
80	<i>M. dissitiflora</i>	A, B
81	<i>M. eleuterostachya</i>	L
82	<i>M. elliptica</i>	A
83	<i>M. ericifolia</i>	A, K
84	<i>M. eulobata</i>	A
85	<i>M. eurystoma</i>	L
86	<i>M. eximia</i>	J
87	<i>M. exuvia</i>	A, L, B/K
88	<i>M. fabri</i>	L
89	<i>M. faucicola</i>	A
90	<i>M. ferruginea</i>	L
91	<i>M. filifolia</i>	A
92	<i>M. fissurata</i>	A
93	<i>M. flammea</i>	A
94	<i>M. flavovirens</i>	A
95	<i>M. fluviatilis</i>	L
96	<i>M. foliolosa</i>	B, L
97	<i>M. formosa</i>	A
98a	<i>M. fulgens</i> subsp. <i>corrugata</i>	A
98b	<i>M. fulgens</i> subsp. <i>fulgens</i>	A
98c	<i>M. fulgens</i> subsp. <i>steedmanii</i>	A
99	<i>M. genialis</i>	J/A
100	<i>M. gibbosa</i>	A
101	<i>M. glaberrima</i>	A
102	<i>M. glauca</i>	A, G
103	<i>M. glena</i>	J
104	<i>M. globifera</i>	A
105	<i>M. glomerata</i>	L/J
106	<i>M. gnidioides</i>	L/J
107	<i>M. grieveana</i>	A
108	<i>M. groveana</i>	G

#	Species	Oil type
109	<i>M. halmaturorum</i>	E
110	<i>M. halophila</i>	A, B
111	<i>M. hamata</i>	A/K, B
112	<i>M. hamulosa</i>	J
113	<i>M. haplantha</i>	A/G/J
114	<i>M. hemisticta</i>	A
115	<i>M. hnatiukii</i>	A
116	<i>M. hollidayi</i>	G
117	<i>M. holosericea</i>	A/J
118	<i>M. howeana</i>	J
119a	<i>M. huegelii</i> subsp. <i>huegelii</i>	J/L
119b	<i>M. huegelii</i> subsp. <i>pristicensis</i>	J/L
120	<i>M. hutensis</i>	K
121	<i>M. hypericifolia</i>	A
122	<i>M. idana</i>	A
123a	<i>M. incana</i> subsp. <i>incana</i>	J,A
123b	<i>M. incana</i> subsp. <i>tenella</i>	J
124	<i>M. interioris</i>	L/J
125	<i>M. irbyana</i>	L/J
126	<i>M. johnsonii</i>	A
127	<i>M. keigheryi</i>	A
128	<i>M. kunzeoides</i>	J
129	<i>M. laetifica</i>	A/J
130	<i>M. lanceolata</i>	A/G/J
131	<i>M. lara</i>	A/G
132	<i>M. lasiandra</i>	G
133	<i>M. lateralis</i>	A
134	<i>M. lateriflora</i>	A
135	<i>M. lateritia</i>	J
136	<i>M. laxiflora</i>	A
137	<i>M. lazaridis</i>	A
138	<i>M. lecanantha</i>	A
139	<i>M. leiocarpa</i>	G
140	<i>M. leiopyxis</i>	K, A, J
141	<i>M. leptospermoides</i>	A
142	<i>M. leucadendra</i>	E, A

#	Species	Oil type
143	<i>M. leuropoma</i>	G
144	<i>M. linariifolia</i>	A, B, C
145	<i>M. linearifolia</i>	A
146a	<i>M. linearis</i> var. <i>linearis</i>	A
146b	<i>M. linearis</i> var. <i>pinifolia</i>	A
147	<i>M. linguiformis</i>	A
148	<i>M. linophylla</i>	A
149	<i>M. longistaminea</i>	J
150	<i>M. lutea</i>	J
151a	<i>M. macronychia</i> subsp. <i>macronychia</i>	A
151b	<i>M. macronychia</i> subsp. <i>trygonoides</i>	A
152	<i>M. manglesii</i>	A
153	<i>M. marginata</i>	J
154	<i>M. megacephala</i>	L
155	<i>M. megalongensis</i>	A
156	<i>M. micromera</i>	A

Oil compositions used in the table

Where percentage figures are mentioned, they are to be taken as only approximate values.

A	50–70% 1,8-cineole, <10% each of α -pinene, limonene, α -terpineol
B	20–50% terpinen-4-ol
C	major amounts of terpinolene
D	lemon-scented oils
E	aromatic oils (in the chemical sense)
F	major amounts of E-nerolidol
G	large amounts of α -pinene
H	significant amounts of β -diketones
J	other sesquiterpenic oils
K	significant amounts of linalool
L	other monoterpenoid oils
M	virtually no oil present
X/Y	oil compositions midway between X and Y type oils
(Z)	isolated occurrences of Z type oils but by no means the major oil type
-	no data
#	species number (notional)

#	Species	Oil type
157	<i>M. microphylla</i>	A
158	<i>M. minutifolia</i>	J
159	<i>M. monantha</i>	G
160	<i>M. montana</i>	J
161	<i>M. montis-zamia</i>	A, A/G
162	<i>M. nanophylla</i>	H
163	<i>M. nematophylla</i>	G
164	<i>M. nervosa</i>	A/K/J
165	<i>M. nesophila</i>	A
166	<i>M. nodosa</i>	A/B
167	<i>M. ochroma</i>	B
168	<i>M. oldfieldii</i>	A
169	<i>M. orbicularis</i>	K
170	<i>M. ordinifolia</i>	A
171	<i>M. orophila</i>	L
172	<i>M. osullivanii</i>	J
173	<i>M. oxyphylla</i>	A
174	<i>M. pachyphylla</i>	L
175	<i>M. pallescens</i>	G
176	<i>M. pallida</i>	L
177	<i>M. paludicola</i>	A
178	<i>M. pancheri</i>	G/J
179	<i>M. papillosa</i>	G/J
180	<i>M. parviceps</i>	K
181	<i>M. parvistaminea</i>	A
182	<i>M. pauciflora</i>	L
183a	<i>M. pauperiflora</i> subsp. <i>fastigata</i>	L
183b	<i>M. pauperiflora</i> subsp. <i>mutica</i>	L
183c	<i>M. pauperiflora</i> subsp. <i>pauperiflora</i>	A
184	<i>M. pearsonii</i>	A
185	<i>M. penicula</i>	J
186a	<i>M. pentagona</i> var. <i>latifolia</i>	L
186b	<i>M. pentagona</i> var. <i>pentagona</i>	L
186c	<i>M. pentagona</i> var. <i>raggedensis</i>	L
187	<i>M. phoenicea</i>	A
188	<i>M. phoidophylla</i>	A/G

#	Species	Oil type
189	<i>M. phratra</i>	M
190	<i>M. pityoides</i>	A
191	<i>M. platycalyx</i>	A
192	<i>M. plumea</i>	A, A/G
193	<i>M. podiocarpa</i>	J
194	<i>M. polandii</i>	M
195	<i>M. polycephala</i>	A/G
196	<i>M. pomphostoma</i>	A/J
197	<i>M. preissiana</i>	J
198	<i>M. pritzelii</i>	G
199	<i>M. procera</i>	L
200	<i>M. protrusa</i>	A
201	<i>M. psammophila</i>	L
202	<i>M. pulchella</i>	A
203	<i>M. pungens</i>	M
204	<i>M. punicea</i>	J/L
205	<i>M. pustulata</i>	A
206	<i>M. pyramidalis</i>	J
207	<i>M. quadrifaria</i>	J
208	<i>M. quericina</i>	M
209	<i>M. quinquenervia</i>	F, A/J/K
210	<i>M. radula</i>	A
211	<i>M. recurva</i>	A
212	<i>M. raphiophylla</i>	A
213	<i>M. rigidifolia</i>	A
214	<i>M. ringens</i>	A
215	<i>M. rugulosa</i>	A
216	<i>M. ryeae</i>	A
217	<i>M. sabrina</i>	L
218	<i>M. salicina</i>	A
219	<i>M. saligna</i>	J/L
220	<i>M. sapientes</i>	A
221	<i>M. scabra</i>	J
222	<i>M. scalena</i>	A
223	<i>M. sciostyloa</i>	-
224	<i>M. sclerophylla</i>	J/L

#	Species	Oil type
225	<i>M. sculponeata</i>	G/J
226	<i>M. seriata</i>	J
227	<i>M. sericea</i>	G
228	<i>M. serpentina</i>	A
229	<i>M. sheathiana</i>	G
230	<i>M. shiressii</i>	A
231	<i>M. sieberi</i>	G
232	<i>M. similis</i>	A
233	<i>M. societatis</i>	J
234	<i>M. sophisma</i>	J
235	<i>M. sparsiflora</i>	A
236	<i>M. spathulata</i>	A/G
237	<i>M. spectabilis</i>	A
238a	<i>M. sphaerodendra</i> var. <i>microphylla</i>	J
238b	<i>M. sphaerodendra</i> var. <i>sphaerodendra</i>	-
239	<i>M. spicigera</i>	A, K
240	<i>M. squamea</i>	J
241	<i>M. squamophloia</i>	E
242	<i>M. squarrosa</i>	L
243	<i>M. stenostachya</i>	A
244	<i>M. stereophloia</i>	A
245	<i>M. stipitata</i>	D
246	<i>M. stramentosa</i>	J
247	<i>M. striata</i>	A
248	<i>M. strobophylla</i>	G
249	<i>M. styphelioides</i>	J
250	<i>M. subalaris</i>	A/B
251	<i>M. suberosa</i>	A
252	<i>M. subfalcata</i>	A
253	<i>M. subtrigona</i>	L
254	<i>M. subulata</i>	A
255	<i>M. sylvana</i>	G
256	<i>M. systema</i>	K
257	<i>M. tamariscina</i>	G
258	<i>M. teretifolia</i>	A, D

#	Species	Oil type
259	<i>M. teuthidoides</i>	A
260	<i>M. thapsina</i>	J/A
261	<i>M. thymifolia</i>	A
262	<i>M. thymoides</i>	M
263	<i>M. thyoides</i>	L, J
264	<i>M. tinkeri</i>	L, J
265	<i>M. torquata</i>	A
266	<i>M. tortifolia</i>	J, M
267	<i>M. trichophylla</i>	A, C
268	<i>M. trichostachya</i>	A, C
269	<i>M. triumphalis</i>	H
270a	<i>M. tuberculata</i> var. <i>arenaria</i>	J/L
270b	<i>M. tuberculata</i> var. <i>macrophylla</i>	L
270c	<i>M. tuberculata</i> var. <i>tuberculata</i>	J, K
271	<i>M. ulicoides</i>	J
272	<i>M. uncinata</i>	A, B

Oil compositions used in the table

Where percentage figures are mentioned, they are to be taken as only approximate values.

A	50–70% 1,8-cineole, <10% each of α -pinene, limonene, α -terpineol
B	20–50% terpinen-4-ol
C	major amounts of terpinolene
D	lemon-scented oils
E	aromatic oils (in the chemical sense)
F	major amounts of E-nerolidol
G	large amounts of α -pinene
H	significant amounts of β -diketones
J	other sesquiterpenic oils
K	significant amounts of linalool
L	other monoterpenoid oils
M	virtually no oil present
X/Y	oil compositions midway between X and Y type oils
(Z)	isolated occurrences of Z type oils but by no means the major oil type
-	no data
#	species number (notional)

#	Species	Oil type
273	<i>M. undulata</i>	A, L
274	<i>M. urceolaris</i>	A/G/K
275	<i>M. uxorum</i>	J
276	<i>M. venusta</i>	J
277	<i>M. villosisepala</i>	L
278a	<i>M. viminalis</i> subsp. <i>rhododendron</i>	-
278b	<i>M. viminalis</i> subsp. <i>viminalis</i>	A, L
279a	<i>M. viminea</i> subsp. <i>appressa</i>	-
279b	<i>M. viminea</i> subsp. <i>demissa</i>	-
279c	<i>M. viminea</i> subsp. <i>viminea</i>	A/J, J
280	<i>M. vinnula</i>	A, J
281	<i>M. violacea</i>	A
282	<i>M. virens</i>	A
283	<i>M. viridiflora</i>	A, L, E
284a	<i>M. williamsii</i> subsp. <i>fletcheri</i>	A
284b	<i>M. williamsii</i> subsp. <i>synoriensis</i>	A
284c	<i>M. williamsii</i> subsp. <i>williamsii</i>	A
285	<i>M. wilsonii</i>	A
286	<i>M. wimmerensis</i>	A
287	<i>M. wonganensis</i>	A
288	<i>M. xerophila</i>	A
289	<i>M. zeteticorum</i>	A
290	<i>M. zonalis</i>	L

Appendix 2 Species with potential for planting for a variety of uses in different climatic zones

The following species of *Melaleuca* show potential for planting and/or trialling on farms, in saline and other land rehabilitation projects etc. and/or are of value (or potential value) as an essential oils source.

(A) Subtemperate climates

<i>Melaleuca</i> species	Shelterbelts, biodiversity corridors, screens, land rehabilitation etc.	Saline soils	Road verges and parks	Lower rainfall areas	Timber (in the round or sawn timber)/ brushwood fencing ^a	Essential oils/ extractives ^b
<i>acuminata</i>	X	X	X	X	Br	
<i>acutifolia</i>	X		X			
<i>alternifolia</i>	X			X		EO(1)
<i>armillaris</i>	X	X	X			
<i>atroviridis</i>	X	X		X	Br	EO(3)
<i>bracteata</i>	X	X	X	X	Ti	Ex(3)
<i>brevifolia</i>	X	X		X		
<i>citrina</i>	X		X			
<i>concreta</i>	X			X	Br	EO(3)
<i>croxfordiae</i>	X		X			
<i>cuticularis</i>	X	X	X			
<i>decora</i>	X		X			
<i>decussata</i>	X	X				
<i>dissitiflora</i>	X			X		EO(1)
<i>elliptica</i>	X			X		

^a Ti = timber; Br = brushwood fencing

^b Essential oils (EO)/extractives (Ex): (1) = plant source for an established industry; (2) = plant source with a high level of potential based on key characteristics; (3) = plant source requiring substantial research and development (growth/oil yield, oil quality/markets) to confirm potential

Note: In a few cases, a species typically, or only, occurs in higher rainfall areas, and typically is planted mainly in such areas. However, the species may also occur in lower rainfall areas and/or be cultivated sparingly in these, in which case its potential for trial and/or broader use in lower rainfall areas is indicated by 'also'.

(A) Subtemperate climates (continued)

<i>Melaleuca</i> species	Shelterbelts, biodiversity corridors, screens, land rehabilitation etc.	Saline soils	Road verges and parks	Lower rainfall areas	Timber (in the round or sawn timber)/ brushwood fencing ^a	Essential oils/ extractives ^b
<i>eleuterostachya</i>	X		X	X		
<i>ericifolia</i>	X	X	X			EO(1)
<i>exuvia</i>	X	X	X	X		EO(3)
<i>glomerata</i>	X	X	X	X		
<i>halmaturorum</i>	X	X	X	X		
<i>halophila</i>		X				EO(3)
<i>hamata</i>	X	X	X	X	Br	EO(3)
<i>hamulosa</i>	X	X		X	Br	
<i>huegelii</i>	X		X	X		
<i>hypericifolia</i>	X					
<i>lanceolata</i>	X	X		also X in NSW, Qld	Ti	
<i>lateriflora</i>	X	X	X	also X		EO(3)
<i>leiocarpa</i>	X			X		
<i>linariifolia</i>	X	X	X			EO(1)
<i>linearis</i>	X					
<i>linguiformis</i>	X	X				
<i>microphylla</i>	X					
<i>nesophila</i>	X	X	X	also X		
<i>pallescens</i>	X		X			
<i>pallida</i>	X		X			
<i>paludicola</i>	X		X			
<i>parvistaminea</i>	X		X			
<i>pauperiflora</i>	X	X		X		
<i>pentagona</i>	X	X	X			
<i>phoenicea</i>	X		X			
<i>preissiana</i>	X	X	X			

^a Ti = timber; Br = brushwood fencing^b Essential oils (EO)/extractives (Ex): (1) = plant source for an established industry; (2) = plant source with a high level of potential based on key characteristics; (3) = plant source requiring substantial research and development (growth/oil yield, oil quality/markets) to confirm potential

Note: In a few cases, a species typically, or only, occurs in higher rainfall areas, and typically is planted mainly in such areas. However, the species may also occur in lower rainfall areas and/or be cultivated sparingly in these, in which case its potential for trial and/or broader use in lower rainfall areas is indicated by 'also'.

(A) Subtemperate climates (continued)

<i>Melaleuca</i> species	Shelterbelts, biodiversity corridors, screens, land rehabilitation etc.	Saline soils	Road verges and parks	Lower rainfall areas	Timber (in the round or sawn timber)/ brushwood fencing ^a	Essential oils/ extractives ^b
<i>quadrifaria</i>		X			X	
<i>rhaphiophylla</i>		X		X		
<i>salicina</i>		X		X		
<i>sieberi</i>		X		X		
<i>squarrosa</i>		X	X	X		
<i>stereophloia</i>		X	X			
<i>strobophylla</i>		X	X	X	X	
<i>styphelioides</i>		X	X	X		Ti
<i>teretifolia</i>		X		X		EO(2)
<i>thyoides</i>		X	X		X	
<i>uncinata</i>		X		X	X	Br
<i>viminalis</i>		X		X	X	EO(3)
<i>viminea</i>		X		X	X	Ti
<i>xerophila</i>		X	X		X	
<i>zeteticorum</i>		X	X		X	

(B) Tropical and subtropical climates

Species	Shelterbelts, biodiversity corridors, screens, land rehabilitation etc.	Saline soils	Road verges and parks	Lower rainfall areas	Timber ^a (in the round or sawn timber)	Essential oils/ extractives ^b
<i>acacioides</i>	X	X			Ti	EO(3)
<i>alsophila</i>	X	X			Ti	EO(3)
<i>alternifolia</i>	X		X			EO(1)
<i>arcana</i>	X		X		Ti	EO(3)
<i>argentea</i>	X		X		Ti	EO(3)
<i>bracteata</i>	X	X	X	X	Ti	Ex(3)
<i>cajuputi</i>	X		X		Ti	EO(1)

^a Ti = timber; Br = brushwood fencing^b Essential oils (EO)/extractives (Ex): (1) = plant source for an established industry; (2) = plant source with a high level of potential based on key characteristics; (3) = plant source requiring substantial research and development (growth/oil yield, oil quality/markets) to confirm potential

Note: In a few cases, a species typically, or only, occurs in higher rainfall areas, and typically is planted mainly in such areas. However, the species may also occur in lower rainfall areas and/or be cultivated sparingly in these, in which case its potential for trial and/or broader use in lower rainfall areas is indicated by 'also'.

(B) Tropical and subtropical climates (continued)

Species	Shelterbelts, biodiversity corridors, screens, land rehabilitation etc.	Saline soils	Road verges and parks	Lower rainfall areas	Timber ^a (in the round or sawn timber)	Essential oils/ extractives ^b
<i>dealbata</i>	X	X	X		Ti	
<i>decora</i>	X		X			
<i>densispicata</i>	X		X	X		
<i>irbyana</i>	X					
<i>lanceolata</i>	X	X		X	Ti	
<i>leucadendra</i>	X	X	X		Ti	
<i>linariifolia</i>	X	X	X			EO(1)
<i>nervosa</i>	X	X	X			
<i>pallescens</i>	X		X	also X		
<i>quinquenervia</i>	X	X	X		Ti	EO(1)
<i>sericea</i>	X		X			
<i>sieberi</i>	X		X			
<i>squamophloia</i>	X					
<i>stenostachya</i>	X				Ti	
<i>stipitata</i>						EO(3)
<i>styphelioides</i>	X	X	X		Ti	
<i>tamariscina</i>	X		X			
<i>trichostachya</i>	X		X			
<i>viminalis</i>	X		X	X		
<i>viminea</i>	X		X			
<i>viridiflora</i>	X		X		Ti	EO(2)

^a Ti = timber^b Essential oils (EO)/extractives (Ex): (1) = plant source for an established industry; (2) = plant source with a high level of potential based on key characteristics; (3) = plant source requiring substantial research and development (growth/oil yield, oil quality/markets) to confirm potential

Note: In a few cases, a species typically, or only, occurs in higher rainfall areas, and typically is planted mainly in such areas. However, the species may also occur in lower rainfall areas and/or be cultivated sparingly in these, in which case its potential for trial and/or broader use in lower rainfall areas is indicated by 'also'.

In addition to the 'Notes' accompanying the 'Species accounts' (Chapter 7) in this publication, the following references were consulted in preparing Appendix 2: Doran and Turnbull (1997); Elliot and Jones (1982, 1993); Marcar and Crawford (2004); Wrigley and Fagg (1993); Holliday (2004).

References

- Aboriginal Communities of the Northern Territory 1993. Traditional Aboriginal medicines in the Northern Territory of Australia. Conservation Commission of the Northern Territory: Darwin.
- Ahmad F.B.H., Hassan V.U., Zakaria R.B. and Ali J.H. 1997. Betulinic and ursolic acid from seed of *Melaleuca cajuputi* (Myrtaceae). Oriental Chemistry 13(3), 231–232.
- Anderson E. 1993. Plants of central Queensland—their identification and uses. Queensland Department of Primary Industries: Brisbane.
- AVONGRO 2007. Brushwood economics. AVONGRO Wheatbelt Tree Cropping Incorporated: Perth.
- Baillon H.E. 1876. *Melaleuca*. Histoire des plantes 6, 359–360.
- Balciunas J.K., Bowman G.J. and Edwards E.D. 1993a. Herbivorous insects associated with the paperbark *Melaleuca quinquenervia* and its allies: I. Noctuoidea (Lepidoptera). Australian Entomologist 20, 13–24.
- Balciunas J.K. and Burrows D.W. 1993. The rapid suppression of the growth of *Melaleuca quinquenervia* saplings in Australia by insects. Journal of Aquatic Plant Management 31, 265–270.
- Balciunas J.K., Burrows D.W. and Edwards E.D. 1993b. Herbivorous insects associated with the paperbark tree *Melaleuca quinquenervia* and its allies: II. Geometridae (Lepidoptera). Australian Entomologist 20, 91–98.
- Bar A., Fatma M., Zaghoul A.M., Bachawal S.V., Sylvester P.W., Ahmad K.F. and El Sayed K.A. 2008. Antiproliferative triterpenes from *Melaleuca ericifolia*. Journal of Natural Products 71(10), 1787–1790.
- Barlow B.A. 1987a, as 1986. *Regelia punicea* (N. Byrnes) Barlow, comb. nov. (Myrtaceae) from the Northern Territory: phytogeographic implications. Brunonia 9, 89–97.
- Barlow B.A. 1987b, as 1986. Contributions to a revision of *Melaleuca* (Myrtaceae): 1–3. Brunonia 9, 163–177.
- Barlow B.A. 1988. Patterns of differentiation in tropical species of *Melaleuca* L. (Myrtaceae). Proceedings of the Ecological Society of Australia 15, 239–247.
- Barlow B.A. and Cowley K.J. 1988. Contributions to a revision of *Melaleuca* (Myrtaceae): 4–6. Australian Systematic Botany 1, 95–126.
- Barlow B.A. and Forrester J. 1984. Pollen tube interaction in *Melaleuca*. Pp. 154–160 in ‘Pollination ’84’, ed. by E.G. Williams and R.B. Knox. School of Botany, University of Melbourne: Melbourne.
- Baskorowati L. 2006. Controlled pollination methods for *Melaleuca alternifolia* (Maiden & Betche) Cheel. ACIAR Technical Reports No. 63. Australian Centre for International Agricultural Research: Canberra.
- Baskorowati L., Doran J.C. and Moncur M.W. 2010a. Reproductive biology of *Melaleuca alternifolia* (Maiden and Betche) Cheel and implications to the breeding system. Lambert Academic Publishing: Saarbrücken, Germany.
- Baskorowati L., Moncur M.W., Cunningham S.A., Doran J.C. and Kanowski P.J. 2010b. Reproductive biology of *Melaleuca alternifolia* (Myrtaceae) 2. Incompatibility and pollen transfer in relation to the breeding system. Australian Journal of Botany 58, 384–391.
- Baskorowati L., Moncur M.W., Doran J.C. and Kanowski P.J. 2010c. Reproductive biology of *Melaleuca alternifolia* (Myrtaceae) 1. Floral biology. Australian Journal of Botany 58, 373–383.
- Bauer K., Garbe D. and Surburg H. 1997. Common fragrance and flavour materials. Wiley-VCH: Weinheim, Germany.
- Bentham G. 1867, as 1866. Myrtaceae. Pp. 1–289 in ‘Flora Australiensis, Volume 3’. Lovell Reeve & Co: London.
- Blake N.M., Wightman G. and Williams L. 1998. Iwaidja ethnobotany. Northern Territory Botanical Bulletin 23, 1–162.
- Blake S.T. 1968. A revision of *Melaleuca leucadendron* and its allies (Myrtaceae). Contributions from the Queensland Herbarium 1, 1–114.
- Blake S.T. and Roff C. 1972. The honey flora of Queensland. Queensland Department of Primary Industries: Brisbane.
- Boland D.J., Brooker M.I.H., Chippindale G.M., Hall N., Hyland B.P.M., Johnson R.D. et al. 2006. Forest trees of Australia, fifth edition. CSIRO Publishing: Melbourne.
- Bootle K.R. 1983. Wood in Australia. McGraw-Hill Book Co.: Sydney.

- Briggs J.D. and Leigh J.H. 1995. Rare and threatened Australian plants. CSIRO Publishing: Melbourne.
- Brighton C.A. and Ferguson I.K. 1976. Chromosome counts in the genus *Melaleuca* (Myrtaceae). Kew Bulletin 31(1), 27–33.
- Broadhurst L., Byrne M., Craven L. and Lepschi B. 2004. Genetic congruence with new species boundaries in the *Melaleuca uncinata* complex (Myrtaceae). Australian Journal of Botany 52(6), 729–737.
- Brock J. 1988. Top End native plants. The author: Winnellie, Darwin.
- Brophy J.J. 1999. Potentially commercial melaleucas. Pp. 247–274 in ‘Tea tree, the genus *Melaleuca*’, ed. by I.A. Southwell and R.F. Lowe. Medicinal and Aromatic Plants: Industrial Profiles, Volume 9. Harwood Academic Publishers: Amsterdam.
- Brophy J.J. and Clarkson J.R. 1989. The essential oils of four chemotypes of *Melaleuca citrolens*. Journal and Proceedings of the Royal Society of New South Wales 122, 11–18.
- Brophy J.J., Craig D.C., Goldsack R.J., Fookes C.J.R., Leach D.N. and Waterman P.G. 2006a. Triumphantone, a novel diketone from the volatile oil of the leaves of *Melaleuca triumphalis*, and its conversion into isotriumphantone. Phytochemistry 67, 2085–2089.
- Brophy J.J. and Doran J.C. 1996. Essential oils of tropical *Asteromyrtus*, *Callistemon* and *Melaleuca* species. ACIAR Monograph No. 40. Australian Centre for International Agricultural Research: Canberra.
- Brophy J.J. and Doran J.C. 2004. Geographic variation in oil characteristics in *Melaleuca ericifolia*. Journal of Essential Oil Research 16, 4–8.
- Brophy J.J., Forster P.I., Goldsack R.J., Hibbert D.B. and Punruckvong A. 1997. Variation in *Callistemon viminalis* (Myrtaceae): new evidence from volatile oils. Australian Systematic Botany 10, 1–13.
- Brophy J.J., Goldsack R.J., Craven L.A., Ford A.J. and Godfree R. 2008. The leaf oils of the *Melaleuca minutifolia* F.Muell. group (Myrtaceae). Journal of Essential Oil Research 20, 163–168.
- Brophy J.J., Goldsack R.J., Craven L.A. and O’Sullivan W. 2006b. An investigation of the leaf oils of the Western Australian broombush complex (*Melaleuca uncinata* sens. lat.) (Myrtaceae). Journal of Essential Oil Research 18, 591–599.
- Brophy J.J., Goldsack R.J., Doran J.C., Craven L.A. and Lepschi B.J. 1999. A comparison of the leaf oils of *Melaleuca squamophloia* with those of its close relatives *M. stypheiosides* and *M. bracteata*. Journal of Essential Oil Research 11, 327–332.
- Brophy J.J., Goldsack R.J., Forster P.I., Craven L.A. and Lepschi B.J. 1998. The leaf essential oils of the Australian members of the genus *Callistemon* (Myrtaceae). Journal of Essential Oil Research 10, 595–606.
- Brophy J.J., Goldsack R.J. and Goldsack G. 1993. The essential oils of the indigenous Myrtaceae of Lord Howe Island. Flavour and Fragrance Journal 8, 153–159.
- Brophy J.J. and Lassak E.V. 1983. The volatile leaf oils of *Melaleuca armillaris*, *M. dissitiflora* and *M. trichostachya*. Journal and Proceedings of the Royal Society of New South Wales 116, 7–10.
- Brophy J.J. and Lassak E.V. 1988. The leaf oil of *Melaleuca leucadendra*. L. Flavour and Fragrance Journal 3, 43–46.
- Brophy J.J. and Lassak E.V. 1992. Steam volatile leaf oils of some *Melaleuca* species from Western Australia. Flavour and Fragrance Journal 7, 27–31.
- Brophy J.J., Lassak E.V. and Boland D.J. 1987. Volatile leaf oils of the two subspecies of *Melaleuca acacioides* F.Muell. Journal and Proceedings of the Royal Society of New South Wales 120, 135–139.
- Brophy J.J., Lassak E.V. and Boland D.J. 1988. Volatile leaf oils of six northern Australian broad-leaved melaleucas. Journal and Proceedings of the Royal Society of New South Wales 112, 29–33.
- Brophy J.J., Lassak E.V. and Boland D.J. 1990. Steam volatile oils of *Melaleuca globifera*, *M. lateriflora*, *M. symphyocarpa* and *M. uncinata*. Flavour and Fragrance Journal 5, 43–48.
- Brophy J.J., Lassak E.V. and Toia R.F. 1985. The volatile leaf oils of *Callistemon viminalis*. Journal and Proceedings of the Royal Society of New South Wales 118, 101–104.
- Brown G.K., Udovicic F. and Ladiges P.Y. 2001. Molecular phylogeny and biogeography of *Melaleuca*, *Callistemon* and related genera (Myrtaceae). Australian Systematic Botany 14, 565–585.
- Brown R. 1814. General remarks, geographical and systematical, on the botany of Terra Australis. Pp. 533–613 (Appendix III) in ‘A voyage to Terra Australis’, Volume 2 by M. Flinders. G. & W. Nicol: London.
- Brundrett M., Bouger N., Dell B., Grove T. and Malajczuk N. 1996. Working with mycorrhizas in forestry and agriculture. ACIAR Monograph No. 32. Australian Centre for International Agricultural Research: Canberra.
- Bulman P., Beale P. and Knight A. 1998. Growing broombush for profit and land protection. Primary Industries and Resources South Australia: Murray Bridge.
- Bultman J.D., Beal R.H., Huffman J.B. and Parrish K.K. 1983. An investigation of the natural resistance of *Melaleuca quinquenervia* to tropical and terrestrial wood-destroying organisms. Forest Products Journal 33, 39–43.
- Burrows D.W., Balciunas J.K. and Edwards E.D. 1994. Herbivorous insects associated with the paperbark *Melaleuca quinquenervia* and its allies: III. Gelechioidea (Lepidoptera). Australian Entomologist 21, 137–142.
- Butcher P.A. 1994. Genetic diversity in *Melaleuca alternifolia*: implications for breeding to improve production of Australian tea tree oil. PhD thesis. Australian National University: Canberra.

- Butcher P.A., Bell J.C. and Moran G.F. 1992. Patterns of genetic diversity and nature of the breeding system in *Melaleuca alternifolia* (Myrtaceae). Australian Journal of Botany 40, 365–375.
- Butcher P.A., Byrne M. and Moran G.F. 1995. Variation within and among the chloroplast genomes of *Melaleuca alternifolia* and *M. linariifolia* (Myrtaceae). Plant Systematics and Evolution 194, 69–81.
- Butcher P.A., Doran J.C. and Slee M.U. 1994. Intraspecific variation in leaf oils of *Melaleuca alternifolia* (Myrtaceae). Biochemical Systematics and Ecology 22, 419–430.
- Byrnes N.B. 1984. A revision of *Melaleuca* L. (Myrtaceae) in northern and eastern Australia, 1. Austrobaileya 2, 65–76.
- Byrnes, N.B. 1985. A revision of *Melaleuca* L. (Myrtaceae) in northern and eastern Australia, 2. Austrobaileya 2, 131–146.
- Byrnes N.B. 1986. A revision of *Melaleuca* L. (Myrtaceae) in northern and eastern Australia, 3. Austrobaileya 2, 254–273.
- CABI 2000. Forestry compendium global module (CD-ROM). Wallingford, United Kingdom: CAB International.
- Cameron L. 2003. Broombush. Department of Primary Industries Agnote AG0638. Victorian Department of Primary Industries: Melbourne.
- Carnegie A.J. and Lidbetter J.R. 2012. Rapidly expanding host range for *Puccinia psidii* sensu lato in Australia. Australasian Plant Pathology 41(1), 13–29.
- Carrick J. and Chorney K. 1979. A review of *Melaleuca* L. (Myrtaceae) in South Australia. Journal of the Adelaide Botanic Gardens 1, 281–319.
- Carter J.L., Colmer T.D. and Veneklaas E.J. 2006. Variable tolerance of wetland tree species to combined salinity and water logging is related to regulation of ion uptake and production of organic solutes. New Phytologist 169(1), 123–133.
- Center T.D. 1992. Biological control of weeds in waterways and on public lands in the southern-eastern United States of America. Pp. 256–263 in 'Proceedings of the First International Weed Control Congress, 17–21 February, 1992, Monash University, Australia', Volume 1, ed. by J.H. Combellack, K.J. Levick, J. Parsons and R.B. Richardson. Weed Science Society of Victoria: Melbourne.
- Chen H.T. and Su Y.C. 1998. Kraft pulping potentials of *Melaleuca leucadendron* wood from Vietnamese plantations. Taiwan Journal of Forest Science 13(3), 209–217.
- Cherrier J.F. 1981. Le niaoulia en Nouvelle-Caledonie (*Melaleuca quinquenervia* S.T. Blake). Revue Forestiere Francaise 33, 297–311.
- Christensen R.M., Pratt P.D., Costello S.L., Rayamajhi M.B. and Center T.D. 2011. Acquired natural enemies of the weed biological control agent *Oxyops vitiosa* (Coleoptera: Curculionidae). Florida Entomologist 94(1), 1–8.
- Clemson A. 1985. Honey and pollen flora. Inkata Press: Melbourne.
- Colton B., Murtagh J., Drinnan J. and Clarke B. 2000. Tea tree oil. Agfact P6.4.6, second edition. New South Wales Agriculture: Orange.
- Cook L.G., Morris D.C., Edwards R.D. and Crisp M.D. 2008. Reticulate evolution in the natural range of the invasive wetland tree species *Melaleuca quinquenervia*. Molecular Phylogenetics and Evolution 47(2), 506–522.
- Coppens J.J.W. 1995. Flavours and fragrances of plant origin. Non-wood Forest Products No. 1. Food and Agriculture Organization of the United Nations (FAO): Rome.
- Courtney J.L., Lassak E.V. and Spiers G.B. 1983. Leaf wax constituents of some myrtaceous species. Phytochemistry 22(4), 947–949.
- Cowley K.J., Quinn F.C., Barlow B.A. and Craven L.A. 1990. Contributions to a revision of *Melaleuca* (Myrtaceae): 7–10. Australian Systematic Botany 3, 165–202.
- Craven L.A. 1989. Reinstatement and revision of *Asteromyrtus* (Myrtaceae). Australian Systematic Botany 1, 373–385.
- Craven L.A. 1999. Generic position of *Melaleuca punicea*: *Petraeomyrtus*, gen. nov. (Myrtaceae). Australian Systematic Botany 12, 675–678.
- Craven L.A. 2006. New combinations in *Melaleuca* for Australian species of *Callistemon* (Myrtaceae). Novon 16, 468–475.
- Craven L.A. 2009. *Melaleuca* (Myrtaceae) from Australia. Novon 19, 444–453.
- Craven L.A. and Dawson J.W. 1998. *Callistemon* of New Caledonia transferred to *Melaleuca* (Myrtaceae). Adansonia, sér. 3 20, 191–194.
- Craven L.A. and Lepisch B.J. 1999. Enumeration of the species and infraspecific taxa of *Melaleuca* (Myrtaceae) occurring in Australia and Tasmania. Australian Systematic Botany 12, 819–927.
- Craven L.A., Lepisch B.J., Broadhurst L. and Byrne M. 2004. Taxonomic revision of the broombush complex in Western Australia (Myrtaceae, *Melaleuca uncinata* s.l.). Australian Systematic Botany 17, 255–271.
- Crowley G., Garnett S. and Shephard S. 2009. Impact of storm-burning on *Melaleuca viridiflora* invasion of grasslands and grassy woodlands on Cape York Peninsula, Australia. Austral Ecology 34(2), 196–209.
- Davis R.L. 2003. The Australian tea tree oil industry. In 'Proceedings of International Federation of Essential Oils and Aroma Trades (IFEAT) International Conference, Sydney, 2–6 November 2003', ed. by C. Green (CD-ROM). IFEAT: London.
- de Oliveira Y., Pinto F., da Silva A.L.L., Guedes I., Biasi L.A. and Quoirin M. 2010. An efficient protocol for micropropagation of *Melaleuca alternifolia* Cheel. In Vitro Cellular and Developmental Biology—Plant 46(2), 192–197.

- Department of Agriculture and Food Western Australia 2004. Salinity tolerance of plants for agriculture and revegetation. At <http://www.agric.wa.gov.au/PC_92359.html>, accessed 23 March 2012.
- Doran J.C. 1990. Nursery practice. Pp. 89–106 in ‘Trees for rural Australia’, ed by K.W. Cremer. Inkata Press: Melbourne.
- Doran J.C. 1997. Seed, nursery practice and establishment. Pp. 59–87 in ‘Australian trees and shrubs: species for land rehabilitation and farm planting in the tropics’, ed. by J.C. Doran and J.W. Turnbull. ACIAR Monograph No. 24. Australian Centre for International Agricultural Research: Canberra.
- Doran J.C. 1999a. Cajuput oil. Pp. 221–233 in ‘Tea tree: the genus *Melaleuca*’, ed. by I.A. Southwell and R.F. Lowe. Medicinal and Aromatic Plants: Industrial Profiles, Volume 9. Harwood Academic Publishers: Amsterdam.
- Doran J.C. 1999b. *Melaleuca cajuputi* Powell. Pp. 126–131 in ‘Plant resources of South-East Asia No. 19: essential oil plants’, ed. by L.P.A Oyen and Nguyen Xuan Dung. Backhuys Publishers: Leiden, The Netherlands.
- Doran J.C., Baker G.R., Williams E.R. and Southwell I.A. 2002. Improving Australian tea tree through selection and breeding (1996–2001). RIRDC Publication No. 07/017. Rural Industries Research and Development Corporation: Canberra.
- Doran J.C., Baker G.R., Williams E.R. and Southwell I.A. 2006. Genetic gains in oil yields after nine years of breeding *Melaleuca alternifolia* (Myrtaceae). Australian Journal of Experimental Agriculture 46, 1521–1527.
- Doran J.C., Drinnan J.E., Macdonell P.F. and Aken K.M. 2007. Screening selected myrtaceous tree species for production of essential oils in northern Queensland. Australian Journal of Experimental Agriculture 47, 755–761.
- Doran J.C. and Turnbull J.W. 1997. Australian trees and shrubs: species for land rehabilitation and farm planting in the tropics. ACIAR Monograph No. 24. Australian Centre for International Agricultural Research: Canberra.
- Dray F.A. Jr, Bennett B.C. and Center T.D. 2006. Invasion history of *Melaleuca quinquenervia* (Cav.) S.T. Blake in Florida. Castanea 71(3), 210–225.
- Edwards R.D., Craven L.A., Crisp M.D. and Cook L.G. 2010. *Melaleuca* revisited: cpDNA and morphological data confirm that *Melaleuca* L. (Myrtaceae) is not monophyletic. Taxon 59, 744–754.
- Eldridge K., Davidson J., Harwood C. and van Wyk G. 1993. Eucalypt domestication and breeding. Clarendon Press: Oxford.
- Elliot H.J., Ohmart C.P. and Wylie F.R. 1998. Insect pests of Australian forests: ecology and management. Inkata Press: Melbourne.
- Elliot W.R. and Jones D.L. 1982. *Callistemon*. Pp. 410–428 in ‘Encyclopaedia of Australian plants suitable for cultivation’, Volume 2. Lothian: Melbourne.
- Elliot W.R. and Jones D.L. 1993. *Melaleuca*. Pp. 315–376 in ‘Encyclopaedia of Australian plants suitable for cultivation’, Volume 6. Lothian: Melbourne.
- El-Toumy S.A.A., Marzouk M.S., Moharram F.A. and Aboutabi E.A. 2001. Flavonoids of *Melaleuca quinquenervia*. Pharmazie 56(1), 94–95.
- Evans J. and Turnbull J.W. 2004. Plantation forestry in the tropics: the role, silviculture, and use of planted forests for industrial, social, environmental, and agroforestry purposes, third edition. Oxford University Press: Oxford.
- Fenton R., Roper R.E. and Watt G.R. 1977. Lowland tropical hardwoods: an annotated bibliography of selected species with plantation potential. External Aid Division, Ministry of Foreign Affairs: Wellington, New Zealand.
- Geary T.F. 1988. *Melaleuca quinquenervia* (Cav.) S.T. Blake. Pp. 567–568 in ‘Useful trees of tropical North America’, ed. by R.M. Burns and M. Mosquera. North American Forestry Commission Publication No. 3. North American Forestry Commission: Washington, DC.
- Geiger R.K. (ed.) 1981. Proceedings of *Melaleuca* symposium, Fort Myers, FL, September 23–24, 1980. Division of Forestry, Florida Department of Agriculture and Consumer Services: Tallahassee.
- Gough D.K., Bell R.E., Ryan P.A. and Bragg C.T. 1989. Drying and burning properties of the wood of some Australian tree species. Pp. 177–186 in ‘Trees for the tropics’, ed. by D.J. Boland. ACIAR Monograph No. 10. Australian Centre for International Agricultural Research: Canberra.
- Gwaze D.P. 1989. Growth and survival of Australian tree species in field trials in Zimbabwe. Pp. 129–138 in ‘Trees for the tropics’, ed. by D.J. Boland. ACIAR Monograph No. 10. Australian Centre for International Agricultural Research: Canberra.
- Habila J.D., Ndukwe G.I., Amupitan J.O., Nok A.J., Mikhail S.A. and Ladan Z. 2010. 3 β -hydroxyllup-20(29)-en-28-oic acid: structural elucidation, *Candidosis* and mould inhibition evaluation. Journal of Medicinal Plants Research 4(18), 1933–1947.
- Hawkeswood T.J. 1980. Jewel beetles as pollinators of *Melaleuca pauperiflora* F.Muell. (Myrtaceae) between Eucla (W.A.) and Koonalda (S.A.). Western Australian Naturalist 14, 238–239.
- Hearne D.A. 1975. Trees for Darwin and northern Australia. Australian Government Publishing Service: Canberra.
- Hellyer R.O. and Lassak E.V. 1968. The steam volatile constituents of *Melaleuca viridiflora*. Australian Journal of Chemistry 21, 2585–2587.
- Hendrati R.L., Baskorowati L. and Kartikawati N. 2002. Reproductive biology of *Melaleuca cajuputi* subsp. *cajuputi*. Pp. 139–143 in ‘Proceedings international seminar—advances in genetic improvement of tropical

- tree species', ed. by A. Rimbawanto and M. Susanto. Centre for Forest Biotechnology and Tree Improvement: Jogjakarta, Indonesia.
- Hnawia E., Brophy J.J., Craven L.A., Lebouvier N., Caballion P. and Nour M. 2012. An examination of the leaf essential oils of the endemic *Melaleuca* (Myrtaceae) species of New Caledonia. *Journal Essential Oil Research* 21, 273–278.
- Hoang Chuong, Doran J.C., Pinyopasarak K. and Harwood C.E. 1996. Variation in growth and survival of *Melaleuca* species in the Mekong Delta of Vietnam. Pp. 31–36 in 'Tree improvement for sustainable tropical forestry', ed. by M.J. Dieters, A.C. Matheson, D.G. Nikles, C.E. Harwood and S.M. Walker. Proceedings of the Queensland Forest Research Institute – International Union of Forest Research Organizations (QFRI–IUFRO) Conference, Caloundra, Queensland, Australia, 27 October – 1 November 1996. Queensland Forestry Research Institute: Gympie.
- Holliday I. 2004. *Melaleucas*: a field and garden guide, second edition. Reed New Holland: Sydney.
- Homer L.E., Leach D.N., Lea D., Lee L.S., Henry R.J. and Baverstock P.R. 2000. Natural variation in the essential oil content of *Melaleuca alternifolia* Cheel (Myrtaceae). *Biochemical Systematics and Ecology* 28, 367–382.
- Huffman J.B. 1981. *Melaleuca* wood and bark utilization research—a progress report. Pp. 37–68 in 'Proceedings of *Melaleuca* symposium, Fort Myers, FL, September 23–24, 1980', ed. by R.K. Geiger. Division of Forestry, Florida Department of Agriculture and Consumer Services: Tallahassee.
- Hussein S.A.M., Hashim A.N.M., El-Sharawy R.T., Seliem M.A., Linscheid U. and Nawwar M.A.M. 2007. Ericifolin: an eugenol 5-O-galloyl glucoside and other phenolics from *Melaleuca ericifolia*. *Phytochemistry* 68(10), 1464–1470.
- Ireland B.F., Hibbert D.B., Goldsack R.J., Doran J.C. and Brophy J.J. 2002. Chemical variation in the leaf oil of *Melaleuca quinquenervia* (Cav.) S.T. Blake. *Biochemical Systematics and Ecology* 30, 457–470.
- ISO (International Organization for Standardization) 2004. Oil of *Melaleuca*, terpinen-4-ol type, 4730. ISO: Geneva.
- James S.H. 1958. Apomixis in the genus *Callistemon* R. Br. MSc thesis. University of Sydney: Sydney. [Copy in CSIRO library, Canberra.]
- Jones D.L. and Elliot W.R. 1986. Pests, diseases and ailments of Australian plants with suggestions for their control. Lothian: Melbourne.
- Kartikawati N.K. 2005. *Tingkat inkompatibilitas bersilang sendiri pada tanaman kayu putih* (The level of self-incompatibility of *Melaleuca cajuputi* subsp. *cajuputi*). *Jurnal Penelitian Hutan Tanaman* 2, 141–147. [In Indonesian, with English summary]
- Keating W.G. and Bolza E. 1982. Characteristics, properties and uses of timbers. South-east Asia, northern Australia and the Pacific, Volume 1. Inkata Press: Melbourne.
- Keszei A., Hassan Y. and Foley W.J. 2010a. A biochemical interpretation of terpene chemotypes in *Melaleuca alternifolia*. *Journal of Chemical Ecology* 36, 652–661.
- Keszei A., Webb H., Külheim C. and Foley W. 2010b. Genetic tools for improving tea tree oils. RIRDC Publication No. 10/189. Rural Industries Research and Development Corporation: Canberra.
- Khan A.G. 1993. Occurrence and importance of mycorrhizae in aquatic trees of New South Wales, Australia. *Mycorrhiza* 3, 31–38.
- Külheim C., Yeoh S.H., Wallis I.R., Laffan S., Moran G.F. and Foley W.J. 2011. The molecular basis of quantitative variation in foliar secondary metabolites in *Eucalyptus globulus*. *New Phytologist* 191, 1041–1053.
- Ladiges P.Y., McFadden G.I., Middleton N., Orlovich D.A., Treloar N. and Udovicic F. 1999. Phylogeny of *Melaleuca*, *Callistemon*, and related genera of the *Beaufortia* suballiance (Myrtaceae) based on 5S and ITS-1 spacer regions of nrDNA. *Cladistics* 15, 151–172.
- Laffan S.W., Lubarsky E. and Rosauer D.F. 2010. Biodiverse, a tool for the spatial analysis of biological and related diversity. *Ecography* 33, 643–647.
- Lamb D. 1975. Kunjingini plantations 1965–1975. Tropical Forestry Research Note Sr. 24. Papua New Guinea Department of Forests: Boroka.
- Lassak E.V. and Brophy J.J. 2004. Steam volatile oil of some Western Australian species of the family Myrtaceae. *Flavour and Fragrance Journal* 19, 12–16.
- Lassak E.V. and McCarthy T. 2011. Australian medicinal plants: a complete guide to identification and usage. Reed New Holland: Sydney.
- Lazarides M., Cowley K. and Hohnen P. 1997. CSIRO handbook of Australian weeds. CSIRO Publishing: Melbourne.
- Lee C.-K. and Chang M.-H. 1999. Four new triterpenes from the heartwood of *Melaleuca leucadendron*. *Journal of Natural Products* 62(7), 1003–1005.
- Levitt D. 1981. Plants and people: aboriginal uses of plants on Groote Eylandt. Australian Institute of Aboriginal Studies: Canberra.
- Lim S.C. and Midon M.S. 2001. Timber of gelam (*Melaleuca cajuputi* Powell). *Timber Technology Bulletin* No. 23. Forest Research Institute Malaysia (FRIM): Kuala Lumpur.
- Linnaeus C. 1754. *Myrtus leucadendra*. Herbarium Amboinense 9.
- Linnaeus C. 1767. *Melaleuca*. *Systema naturae* ed. 12, 2, 509; *Mantissa plantarum* 1: 14, 105, *nom. cons.*
- Loope L., Duever M., Herndon A., Snyder J. and Jansen D. 1994. Hurricane impact on uplands and freshwater swamp forest. *BioScience* 44, 238–246.

- Lowry J.B. 1973. A new constituent of biogenetic, pharmacological and historical interest from *M. cajeputi* oil. *Nature* 241, 61–62.
- Lum S.K.Y. 1994. Dispersal of Australian plants across Wallace's Line: a case study of *Melaleuca cajuputi* (Myrtaceae). The author: Berkeley.
- McJannet D. 2008. Water table and transpiration dynamics in a seasonally inundated *Melaleuca quinquenervia* forest, north Queensland. *Hydrological Processes* 22(16), 3079–3090.
- McKelvie L., Bills J. and Peat A. 1994. Jojoba, blue mallee and broombush: market assessment and outlook. ABARE Research Report 94.9. Australian Bureau of Agricultural and Resource Economics: Canberra.
- McNeill J., Barrie F.R., Buck W.R., Demoulin V., Greuter W., Hawksworth D.L. et al. 2012. International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. *Regnum Vegetabile* 154. Koeltz: Königstein. Accessible at <<http://www.iapt-taxon.org/nomen/main.php>>.
- Maiden J.H. 1889. The useful native plants of Australia. Compendium: Melbourne.
- Marcar N. and Crawford D. 2004. Trees for saline landscapes. RIRDC Publication No. 03/108. Rural Industries Research and Development Corporation: Canberra.
- Marcar N.E., Crawford D.F., Leppert P.L., Jovanovic T., Floyd R. and Farrow R. 1995. Trees for saltland: a guide to selecting native species for Australia. CSIRO Publishing: Melbourne.
- Martin M.R., Tipping P.W., Reddy K.R., Madeira P.T. and Fitzgerald D. 2011. An evaluation of the impact of *Melaleuca quinquenervia* invasion and management on plant community structure after fire. *Aquatic Botany* 95(4), 287–291.
- Masunga P. 1998 Analysis of the essential oils of *Melaleuca bracteata* from eastern Australia. M. Chem. thesis. University of New South Wales: Sydney.
- Mensforth L.J. and Walker G.R. 1996. Root dynamics of *Melaleuca halmaturorum* in response to fluctuating saline groundwater. *Plant and Soil* 184(1), 75–84.
- Molyneux W. 1993. A new species of *Callistemon* R.Br. (Myrtaceae) from East Gippsland. *Muelleria* 8, 61–64.
- Molyneux W. 1994. *Callistemon kenmorrisonii* (Myrtaceae), a new species from East Gippsland. *Muelleria* 8, 379–383.
- Molyneux W. 1997. Notes on *Callistemon* in East Gippsland, including the description of *C. genofluvialis* sp. nov. *Muelleria* 10, 57–61.
- Molyneux W.M. 2005. *Callistemon nyallengensis* (Myrtaceae) a new species from East Gippsland, Victoria. *Muelleria* 21, 101–104.
- Morin L., Aveyard R., Lidbetter J.R. and Wilson P.G. 2012. Investigating the host-range of the rust fungus *Puccinia psidii* sensu lato across tribes of the family Myrtaceae present in Australia. *PLoS One* 7(4), e35434. At <<http://www.plosone.org/article/info:doi/10.1371/journal.pone.0035434>>, accessed 23 February 2013.
- Morton J.F. 1966. The cajeput tree—a boon and an affliction. *Economic Botany* 20, 31–39.
- Mueller F. 1864. *Callistemon salignus*. *Fragmenta photographiae Australiae* 4, 54–55.
- Mulder V. 1992. Honey and wax production in submerged *Melaleuca* forests in Vietnam. *Bos Newsletter* No. 26, 11(2), 15–29.
- Naidu B.P. 2003. Production of betaine from Australian *Melaleuca* spp. for use in agriculture to reduce plant stress. *Australian Journal of Experimental Agriculture* 43, 1163–1170.
- Naidu B.P. and Cameron D.F. 1999. Reducing plant stress using Australian melaleuca: a report for the Rural Industries Research and Development Corporation. RIRDC Publication No. 99/148. Rural Industries Research and Development Corporation: Canberra.
- Naidu B.P., Paleg L.G. and Jones G.P. 2000. Accumulation of proline analogues and adaptation of *Melaleuca* species to diverse environments in Australia. *Australian Journal of Botany* 48(5), 611–620.
- NAS (National Academy of Sciences) 1983. Firewood crops: shrub and tree species for energy production, Volume 2. National Academic Press: Washington, DC.
- Nguyen N.T., Saneoka H., Suwa R. and Fujita K. 2009. The interaction among provenances of *Melaleuca leucadendra* (weeping paperbark), salt, and aluminum. *Forest Science* 55(5), 443–454.
- Padovan I., Keszei A., Kolliner T.G., Degenhardt J. and Foley W.J. 2010. The molecular basis of host plant selection in *Melaleuca quinquenervia* by a successful biological control agent. *Phytochemistry* 71, 1237–1244.
- Pannell D.J. and Ewing M.A. 2004. Managing secondary dryland salinity: options and challenges. In 'New directions for a diverse planet: Proceedings of the Fourth International Crop Science Congress, 26 Sep – 1 Oct 2004, Brisbane, Australia' (CD-ROM). The Regional Institute Ltd: Gosford. Accessible at <www.cropscience.org.au/icsc2004/>.
- Penfold A.R. and Grant R. 1925. The germicidal values of some Australian essential oils and their pure constituents, together with those of some essential oil isolates and synthetics. Part 3. *Journal and Proceedings of the Royal Society of New South Wales* 58, 346–350.
- Pinyopasarak K. 1989. Growth and survival of Australian tree species in field trials in Thailand. Pp. 109–127 in 'Trees for the tropics', ed. by D.J. Boland. ACIAR Monograph No. 10. Australian Centre for International Agricultural Research: Canberra.
- Prastyono, Doran J.C., Nichols J.D. and Raymond C.A. 2011. Potential of clones in improving the financial

- benefits of essential oil production from *Melaleuca alternifolia* plantations. Agroforestry Systems 83(2), 257–266.
- Pratt P.D. and Arakelian G. 2011. First report of the biological control agent *Boreioglycaspis melaleucae* (Hemiptera: Psyllidae) in California, USA. Florida Entomologist 94(3), 721–722.
- Pujiarti R., Ohtani Y. and Ichiumura H. 2011. Physiochemical properties and chemical compositions of *Melaleuca leucadendron* leaf oils taken from plantations in Java, Indonesia. Journal of Wood Science 57(5), 446–451.
- Purcell M.F. and Balciunas J.K. 1994. Life history and distribution of the Australian weevil *Oxyops vitiosa* (Coleoptera: Curculionidae), a potential biological control agent for *Melaleuca quinquenervia* (Myrtaceae). Annals of the Entomological Society of America 87, 867–873.
- Puruntatameri J., Puruntatameri R., Puruntatameri A., Burak L., Tipuamantymirri C., Tipakalippa M. et al. 2001. Tiwi plants and animals. Northern Territory Botanical Bulletin 24, 1–191.
- Ramanoelina P.A.R., Bianchini J.P. and Gaydou E.M. 2008. Main industrial niaouli (*Melaleuca quinquenervia*) oil chemotype productions from Madagascar. Journal of Essential Oil Research 20(3), 261–266.
- Randall R.P. 2002. A global compendium of weeds. R.G. and F.J. Richardson: Meredith, Victoria.
- Rayachhetry M.B., Blakeslee G.M. and Center T.D. 1996a. Predisposition of melaleuca (*Melaleuca quinquenervia*) to invasion by the potential biological control agent *Botryosphaeria ribis*. Weed Science 44, 603–608.
- Rayachhetry M.B., Blakeslee G.M. and Charudattan R. 1996b. Susceptibility of *Melaleuca quinquenervia* to *Botryosphaeria ribis*, a potential biological control agent. Plant Disease 80, 145–150.
- Rayachhetry M.B., Van T.K., Center T.D. and Elliott M.L. 2001. Host range of *Puccinia psidii*, a potential biological control agent of *Melaleuca quinquenervia* in Florida. Biological Control 22(1), 38–45.
- Rayamajhi M.B., Pratt P.D., Center T.D., Tipping P.W. and Van T.K. 2009. Decline in exotic tree density facilitates increased plant diversity: the experience from *Melaleuca quinquenervia* invaded wetlands. Wetlands Ecology and Management 17(5), 455–467.
- Rayamajhi M.B., Pratt P.D., Center T.D. and Van T.K. 2010a. Insects and a pathogen suppress *Melaleuca quinquenervia* cut-stump regrowth in Florida. Biological Control 53(1), 1–8.
- Rayamajhi M.B., Pratt P.D., Center T.D. and Wheeler G.S. 2010b. Differential response by *Melaleuca quinquenervia* trees to attack by the rust fungus *Puccinia psidii* in Florida. Plant Disease 94(8), 1165.2
- Rayamajhi M.B., Van T.K., Center T.D., Goolsby J.A., Pratt P.D. and Racelis A. 2002. Biological attributes of the canopy-held *Melaleuca* seeds in Australia and Florida, US. Journal of Aquatic Plant Management 40, 87–91.
- Richardson F.J., Richardson R.G. and Shepherd R.C.H. 2011. Weeds of the south-east: an identification guide for Australia, second edition. R.G. and F.J. Richardson: Meredith, Victoria.
- RIRDC (Rural Industries Research and Development Corporation) 2007. The effectiveness and safety of Australian tea tree oil. RIRDC Publication No. 07/143. RIRDC: Canberra.
- Robinson C. and Emmott T. no date. Growing broombush: for fencing products on cleared farmland in southern WA. Land Care Program, Department of Agriculture, Fisheries and Forestry: Canberra.
- Robinson F.A. 1981. Relationship of *Melaleuca* to bee-keeping. Pp. 79–80 in ‘Proceedings of Melaleuca Symposium, Fort Myers, FL, September 23–24, 1980’, ed. by R.K. Geiger. Division of Forestry, Florida Department of Agriculture and Consumer Services: Tallahassee.
- Robinson R.W., James E.A. and Boon P.I. 2012. Population structure in the clonal, woody wetland plant *Melaleuca ericifolia* (Myrtaceae): an analysis using historical aerial photographs and molecular techniques. Australian Journal of Botany 60, 9–19.
- Roff C. 1966. Honey flora of north Queensland. Queensland Agricultural Journal 66, 99–105.
- Rosseto M., Slade R.W., Baverstock P.R., Henry R.J. and Slee L.S. 1999. Microsatellite variation and assessment of genetic structure in tea tree (*Melaleuca alternifolia* – Myrtaceae). Molecular Ecology 8, 633–643.
- Rumphius G.E. 2011 (originally published in 1741, here translated, annotated, and with an introduction by E.M. Beekman). The Amboinese herbal: being a description of the most noteworthy trees, shrubs, herbs, land- and water-plants which are found in Amboina and the surrounding islands according to their shape, various names, cultivation, and use : together with several insects and animals : for the most part with the figures pertaining to them : all gathered with much trouble and diligence over many years and described in twelve books, Volumes 1–6. Yale University Press: New Haven, Connecticut.
- Ryan P.A. and Bell R.E. 1989. Growth, coppicing and flowering of Australian tree species in trials in southeast Queensland, Australia. Pp. 49–68 in ‘Trees for the tropics’, ed. by D.J. Boland. ACIAR Monograph No. 10. Australian Centre for International Agricultural Research: Canberra.
- Rye B.L. 1979. Chromosome number variation in the Myrtaceae and its taxonomic implications. Australian Journal of Botany 27, 547–573.
- Safford R.J., Maltby E., Duong Van Ni and Branch N.P. 2009. *Melaleuca* wetlands and sustainable development in the Mekong Delta, Vietnam. Pp. 829–846 in ‘The

- wetlands handbook, first edition', ed. by E. Maltby and T. Barker. Blackwell Publishing: Oxford.
- Sell C. 2010. Chemistry of essential oils. Pp. 121–150 in 'Handbook of essential oils—science, technology and applications', ed. by K.H.C. Baser and G. Buchbauer. CRC Press: Boca Raton, Florida.
- Serbesoff-King K. 2003. *Melaleuca* in Florida: a literature review on taxonomy, distribution, biology, ecology, economic importance and control measures. *Journal of Aquatic Plant Management* 41, 98–112.
- Simpson J. 1995. The *Melaleuca* research program and results to date. Paper in Proceedings of the National Technical Workshop, Forest Based Development of Long Xuyen Quadrangle, 3–5 August 1995, Long Xuyen, An Giang, Vietnam.
- Southwell I.A. 1999. Tea tree constituents. Pp. 29–62 in 'Tea tree, the genus *Melaleuca*', ed. by I.A. Southwell and R.F. Lowe. Medicinal and Aromatic Plants: Industrial Profiles, Volume 9. Harwood Academic Publishers: Amsterdam.
- Southwell I.A. and Lowe R.F. 1999. Tea tree: the genus *Melaleuca*. Medicinal and Aromatic Plants: Industrial Profiles, Volume 9. Harwood Academic Publishers: Amsterdam.
- Southwell I.A., Russell M., Smith R.L., Brophy J.J. and Day J. 2003. *Melaleuca teretifolia* chemovars: new Australian sources of citral and 1,8-cineole. *Journal of Essential Oil Research* 15, 339–341.
- Southwell I.A., Russell M., Smith R.L., Brophy J.J. and Day J. 2005. *Melaleuca teretifolia*, a novel aromatic and medicinal plant from Australia. *Acta Horticulturae* 677, 79–83.
- Southwell I.A., Stiff I.A. and Brophy J.J. 1992. Terpinolene varieties of *Melaleuca*. *Journal of Essential Oil Research* 4, 363–367.
- Stablein J.J., Bucholtz G.A. and Lockey R.F. 2002. Melaleuca tree and respiratory disease. *Annals of Allergy, Asthma & Immunology* 89, 523–530.
- Stewart, A. 2012. Creating an Australian garden. Allen & Unwin: Sydney.
- Streets R.J. 1962. Exotic forest trees in the British Commonwealth. Clarendon Press: Oxford and London.
- Sun D. and Dickinson G.R. 1995. Survival and growth responses of a number of Australian tree species planted on a saline site in tropical north Australia. *Journal of Applied Ecology* 32, 817–826.
- Susanto M., Doran J., Arnold R. and Rimbawanto A. 2003. Genetic variation in growth and oil characteristic of *Melaleuca cajuputi* subsp. *cajuputi* and potential for genetic improvement. *Journal of Tropical Forest Science* 15(3), 469–482.
- Susanto M., Rimbawanto A., Prastyono and Kartikawati N.K. 2010. Genetic improvement in the breeding of *Melaleuca cajuputi* subsp. *cajuputi*. *Jurnal Penelitian Hutan Tanaman* 2(2), 231–241. [In Indonesian, with English summary]
- Sweedman L. 2006. Australian seeds: a photographic guide. Pp. 67–172 in 'Australian seeds: a guide to their collection, identification and biology', ed. by L. Sweedman and D. Merritt. CSIRO Publishing: Melbourne.
- Tanaka K., Masumori M., Yamanoshita T. and Tange T. 2011. Morphological and anatomical changes of *Melaleuca cajuputi* under submergence. *Trees—Structure and Function* 25(4), 695–704.
- The Paperbark Co. 2013. Our products. The Paperbark Co.: Harvey, Western Australia. At <http://www.paperbarkoils.com.au/essential-oils.html#body_mist_honeymyrtle>, accessed 11 May 2013.
- Thornhill A.H., Hope G.S., Craven L.A. and Crisp M.D. 2012. Pollen morphology of the Myrtaceae. Part 2: tribes Backhousieae, Melaleuceae, Metrosidereae, Osborniae and Syzygiae. *Australian Journal of Botany* 60, 200–224.
- Tipping P.W., Martin M.R., Nimmo K.R., Pierce R.M., Smart M.D., White E., Madeira P.T. and Center T.D. 2009. Invasion of a West Everglades wetland by *Melaleuca quinquenervia* countered by classical biological control. *Biological Control* 48(1), 73–78.
- Trilles B., Bouraïma-Madjebi S. and Valet G. 1999. *Melaleuca quinquenervia* (Cavanilles) S.T.Blaire, niaouli. Pp. 237–247 in 'Tea tree, the genus *Melaleuca*', ed. by I.A. Southwell and R.F. Lowe. Medicinal and Aromatic Plants: Industrial Profiles, Volume 9. Harwood Academic Publishers: Amsterdam.
- Trilles B.L., Bombarda I., Bouraïma-Madjebi S., Raharivelomanana P., Bianchini J.-P. and Gaydou E.M. 2006. Occurrence of various chemotypes in niaouli [*Melaleuca quinquenervia* (Cav.) S.T. Blake] essential oil from New Caledonia. *Flavour and Fragrance Journal* 21, 677–682.
- Turnbull J.W. 1986. Multipurpose Australian trees and shrubs; lesser-known species for fuelwood and agroforestry. ACIAR Monograph No. 1. Australian Centre for International Agricultural Research: Canberra.
- Turner C.E., Center T.D., Burrows D.W. and Buckingham G.R. 1998. Ecology and management of *Melaleuca quinquenervia*, an invader of wetlands in Florida, U.S.A. *Wetlands Ecology and Management* 5, 165–178.
- Venning J. 1988. Growing trees for farms, parks and roadsides: a revegetation manual. Lothian Publishing Company: Melbourne.
- Vieira T.R., Barbosa L.C.A., Maltha C.R.A., Paula V.F. and Nascimento E.A. 2004. Chemical constituents from *Melaleuca alternifolia*. *Química Nova* 27(4), 536–539.
- von Carlowitz P.G. 1991. Multipurpose trees and shrubs: sources of seeds and inoculants. International Council for Research in Agroforestry (ICRAF): Nairobi, Kenya.
- Wang S., Huffman J.B. and Littell R.C. 1981. Characterization of *Melaleuca* biomass as fuel for direct combustion. *Wood Science* 13, 216–219.

- Wang S., Huffman J.B. and Rockwood D.L. 1982. Qualitative evaluation of fuelwood in Florida—a summary report. *Economic Botany* 36(4), 381–388.
- Wang S. and Littell R.C. 1983. Phenotypic variation in calorific value of *Melaleuca* materials from south Florida. *Economic Botany* 37, 292–298.
- Webb H.O., Lanfear R., Hamill J., Foley W.J. and Kühlheim C. 2013. The yield of essential oils in *Melaleuca alternifolia* (Myrtaceae) is regulated through transcript abundance of genes in the MEP pathway. *PLoS One* 8(3), e60631. At <<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0060631>>, accessed 25 April 2013.
- Wiersema J.H. and León B. 2013. World economic plants, second edition. CRC Press: Boca Raton.
- Williams C. 2011. Medicinal plants in Australia, Volume 2: gums, resins, tannin and essential oils. Rosenberg Publishing: Sydney.
- Wiynjorrotj P., Flora S. (deceased), Brown N.D. (deceased), Jatbula P. (deceased), Galmurr J., Katherine M. et al. 2005. Jawoyn plants and animals: Aboriginal flora and fauna knowledge from Nitmiluk National Park and the Katherine area, northern Australia. *Northern Territory Botanical Bulletin* 29, 1–234.
- Wollenweber E., Wehde R., Dorr M., Lang G. and Stevens J.F. 2000. C-methyl-flavonoids from the leaf waxes of some Myrtaceae. *Phytochemistry* 55(8), 965–970.
- Wrigley J.W. and Fagg M. 1993. Bottlebrushes, paperbarks and tea trees, and all other species in the *Leptospermum* alliance. Angus and Robertson: Sydney.
- Wrigley J.W. and Fagg M. 2007. Australian native plants: cultivation, use in landscaping and propagation, concise edition. Reed New Holland: Sydney.
- Yaacob K.B., Abdullah C.M. and Joulain D. 1989. Essential oil of *Melaleuca cajuputi*. P. 140 in 'Proceedings of the 11th International Congress of Oils, Fragrances and Flavours, New Delhi, India, 12–16th November 1989', Volume 1, ed. by S.C. Bhattacharyya, N. Sen and K.L. Sethi. Oxford & IBH Publishing: New Delhi.
- Yao L., Jiang Y., Singanusong R., D'Arcy B., Datta N. and Raymont K. 2004. Flavonoids in Australian *Melaleuca*, *Guioa*, *Lophostemon*, *Banksia* and *Helianthus* honeys and their potential for floral authentication. *Food Research International* 37(2), 166–174.
- Yoshimura M., Ito H., Miyashita K., Hatano T., Taniguchi S., Amakura Y. and Yoshida T. 2008. Flavonol glucuronides and C-glucosidic ellagitannins from *Melaleuca squarrosa*. *Phytochemistry* 69(18), 3062–3069.
- Yunupingu B., Yunupingu-Marika L., Marika D., Marika B., Marika B., Marika R. and Wightman G. 1995. Rirratjingu ethnobotany: Aboriginal plant use from Yirrkala, Arnhem Land, Australia. *Northern Territory Botanical Bulletin* 21, 1–112.
- Zhang G.J., Ivkovic M., Prastyono and Doran J.C. 2011. Development of an economic selection index for Australian tea tree. *Agroforestry Systems* 82, 51–60.